

# Endemic Kaposi's Sarcoma in Human Immunodeficiency Virus Type 1-Seronegative Persons: Demonstration of Retrovirus-Like Particles in Cutaneous Lesions

Klemens Rappersberger, Erwin Tschachler, Eva Zonzits, Reinhard Gillitzer, Angelos Hatzakis, Andreas Kaloterakis, Dean L. Mann, Theresa Popow-Kraupp, Robert J. Biggar, Rudolf Berger, Joannis Stratigos, Klaus Wolff, and Georg Stingl

Division of Cutaneous Immunobiology, Department of Dermatology I (KR, ET, EZ, RG, RB, KW, GS) and the Institute of Virology (TP-K), University of Vienna Medical School, Vienna, Austria; the Departments of Hygiene and Epidemiology (AH), Internal Medicine (AK), and Dermatology (JS), University of Athens Medical School, Athens, Greece; the Laboratory of Viral Carcinogenesis (DLM) and the Viral Epidemiology Section (RJB), National Cancer Institute, Frederick and Bethesda, Maryland, U.S.A.

In 1984, Greek physicians reported on the clustering of cases of Kaposi's sarcoma (KS) on the Peloponnesus peninsula. To gain more insight into its pathogenesis, we studied the sero-epidemiologic and clinicopathologic characteristics of 12 Greek KS patients (eight male/four female) five of whom were residents of an endemic area on the Peloponnesus. These patients were in good general health with ages ranging from 48 to 80 years, had no clinical signs of immunodeficiency, and combined the features of both classic and epidemic KS in that they displayed not only involvement of acral areas but also widespread mucocutaneous lesions. Routine laboratory data were within normal limits; no patient had HTLV-1 and HIV-1/2 antibodies, but all patients had

antibodies to several herpesviruses. The histopathology was characteristic of KS with the peculiar feature of a dense infiltrate composed predominantly of CD4<sup>+</sup> T lymphocytes. Immunoenzymatic/morphologic studies of the KS cells were consistent with their origin from lymphatic endothelium. Outstanding ultrastructural findings were tubuloreticular structures and cylindrical confronting cisternae, structures that are indicative of an ongoing viral infection. Indeed, extensive electronmicroscopic studies resulted in the detection of retrovirus-like particles in close association to KS cells in five of 12 patients. This in situ observation opens the possibility that this retro-virus contributes to KS development. *J Invest Dermatol* 95:371-381, 1990

**C**linicopathologically, four distinctive forms of Kaposi's sarcoma (KS) are recognized: (a) the classical variant, which frequently occurs among elderly men of Jewish Ashkenazi ancestry and usually runs a rather benign course over years [1,2]; (b) the endemic African form, which involves children, adolescents, and adults with a high frequency of extracutaneous manifestations and a more ag-

gressive course [3,4]; (c) KS seen in organ transplant recipients undergoing immunosuppressive therapy [5-7]; (d) epidemic KS representing the most important opportunistic neoplasm occurring in human immunodeficiency virus type 1 (HIV-1)-infected individuals [8-12]. The pathogenic mechanisms leading to the histopathologically uniform but clinically diverse KS variants are still unclear. Although immunosuppression apparently contributes to the devel-

Manuscript received May 31, 1990; accepted for publication June 28, 1990.

This work was supported, in part, by an Erwin Schrödinger fellowship (KR, grant number 0426-MED) from the Austrian Science Foundation, Vienna, Austria; by a grant from Schering AG, Berlin; and by a grant from the Deutsche Forschungsgemeinschaft (RG, FRG).

Reprint requests to: Dr. Georg Stingl, Division of Cutaneous Immunobiology, Department of Dermatology I, University of Vienna Medical School, Alser Strasse 4, A-1090 Vienna, Austria.

## Abbreviations:

AIDS: acquired immunodeficiency syndrome  
CCC: cylindrical confronting cisternae  
CMV: cytomegalovirus  
EA: EBV early antigen  
EBNA: EBV nuclear antigen  
EBV: Epstein-Barr virus  
ELISA: enzyme-linked immunosorbent assay

HHV-6: human herpes virus type 6  
HIV-1,2: human immunodeficiency virus types 1 or 2  
HSV: herpes simplex virus  
HTLV-1: human T-lymphotropic virus 1  
IF: immunofluorescence  
Ig: immunoglobulin(s)  
KS: Kaposi's sarcoma  
MoAb: monoclonal antibody(ies)  
PBS: phosphate-buffered saline  
RT: room temperature  
TPHA: treponema pallidum hemagglutination test  
TRS: tubulo-reticular structures  
VCA: EBV capsid antigen  
VDRL: venereal disease research laboratory test  
vWF: von Willebrand factor  
VZV: varicella zoster virus





**Figure 1.** Clinical features of mucocutaneous KS lesions in Greek patients. *a, b*, involvement of the facial skin by multiple KS lesions in patch and plaque stage (*a*); prominent, plaque-like, and nodular lesions on the left ear (*b*). *c–e*, disseminated, streaky, and coin-sized lesions on the extremities. *f*, KS lesions in all stages involving both feet. *g, h*, mucosal KS lesions involving the glans penis (*g*) and the hard palate (*h*).




**Table I.** List of First-Step Antibody Reagents

Reagents <sup>a</sup>	Working Dilution	Reactivity on Normal Human Tissue	Source <sup>b</sup>	Relevant References
PAL-E	1:500	Endothelial cells of mature blood vessels; dermo-epidermal junction	Serotec	[20,21]
EN-4	1:500	Endothelial cells of mature blood and lymphatic vessels	Serotec	[21,22]
Anti-von Willebrand Factor	1:500	Endothelial cells of mature blood vessels	Dakopatts	[23,24]
OKM-5 (anti-CD36)	1:300	Endothelial cells of mature blood vessels; subset of tissue macrophages	Ortho	[24,25]
Anti-laminin	1:500	Basement membrane zone protein	Serotec	[26,27]
Anti-collagen type IV	1:2000	Basement membrane zone protein	Serotec	[26,27]
Anti-desmin	1:100	Intermediate-sized filaments of smooth muscle cells	Dakopatts	[28]
Anti-vimentin	1:50	Intermediate-sized filaments of certain mesenchymal (e.g., endothelial cells, fibroblasts, macrophages) and neuroectodermal (e.g., melanocytes, Schwann cells) cells	Dakopatts	[29]
Anti-leucocyte (HLe-1) (anti-CD45)	1:200	All human leucocytes	Becton-Dickinson	[30]
Anti-Leu-4 (anti-CD3)	1:600	T-cell antigen associated with T-cell receptor	Becton-Dickinson	[31]
Anti-Leu-3a (anti-CD4)	1:200	Human T helper/inducer cell antigen	Becton-Dickinson	[31]
Anti-Leu-2a (anti-CD8)	1:150	Human T cytotoxic/suppressor cell antigen	Becton-Dickinson	[31]
Anti-Leu-M3 (anti-CD14)	1:100	A human monocyte/macrophage antigen	Becton-Dickinson	[32]
B1 (anti-CD20)	1:200	A human B lymphocyte antigen	Coulter	[33]
Anti-Factor XIIIa	1:20,000	Factor XIIIa expressed on mononuclear phagocytes including dermal dendrocytes	Calbiochem	[34]

<sup>a</sup> All reagents were monoclonal antibodies (MoAb) with the exception of the polyclonal rabbit-anti human Factor XIIIa antiserum.

<sup>b</sup> Serotec, Oxford, England, UK; Dakopatts, Copenhagen, DK; Ortho Diagnostic Systems, Raritan, NY, USA; Becton-Dickinson Immunocytometric Systems, Mountain View, CA, USA; Coulter Immunology, Hialeah, FL, USA; Calbiochem Corporation, San Diego, CA, USA.



**Figure 2.** Histopathology of KS lesions of Greek patients. *a*, angiomatous KS: note the irregular size and shape of vascular lumina, outlined by a monolayer endothelium. *b*, sarcomatous KS: spindle cells forming solid tumor fascicles; red blood cell extravasation between the tumor cells.



**Table II.** Mucocutaneous Localization of KS Lesions in Greek Patients\*

Patient	Feet	Thighs	Hands	Arms	Head/Face	Trunk	Oral/Genital Mucosa	Gastrointestinal Tract		
								Esophagus	Stomach	Duodenum
1	+	+	+	+	+	—	—/—	—	++	—
2	++	+	++	—	—	—	—/—	n.d.	n.d.	n.d.
3	++	++	++	+	++	—	+/+	—	++	—
4	++	+	—	—	—	—	—/—	—	++	—
5	+	+	++	+	—	—	—/—	—	++	—
6	+	—	++	—	+	+	+/-	—	++	+
7	++	—	+	+	—	—	+/+	—	—	—
8	++	—	+	+	—	—	—/—	—	+	+
9	—	+	—	—	+	—	—/—	+	+	+
10	—	—	—	++	—	—	+/+	—	++	—
11	++	++	—	—	—	—	—/—	—	++	++
12	+	++	+	++	—	—	—/—	n.d.	n.d.	n.d.

\* Grade of involvement by KS: +, moderate; ++, severe; —, not involved; n.d., not explored by endoscopy.

opment of KS in transplant recipients [7], certain epidemiologic aspects of KS [3] as well as the lack of correlation between severity of immunosuppression and frequency of KS in some HIV-1-infected persons [2] essentially exclude the possibility that immunosuppression is the only factor that triggers the events leading to this proliferative disorder.

In 1984, Kaloterakis reported a clustering of KS among the inhabitants of distinct areas on the Southern Peloponnesus peninsula in Greece [13]. In 1988, an international study group was formed in order to conduct more extensive epidemiologic and clinicopathologic investigations on this cluster of patients and, by doing so, to contribute to the elucidation of pathomechanisms operative in KS.

#### PATIENTS AND METHODS

**Epidemiology** Reportedly, the overall incidence of KS in Greece is 3.5/10<sup>6</sup> people/year with an endemic focus on the Peloponnesus (7.7 cases/10<sup>6</sup> people/year) (Hatzakis et al, in preparation). Twelve KS patients, eight male and four female, consecutively admitted to the Department of Dermatology, Athens University Medical School, were selected for a thorough medical examination. Their ages ranged from 48 to 80 years (mean, 72.6 years) with an average disease duration of 2.5 years (range, 1–5 years). Five patients were born on the Peloponnesus (patients 3, 5, 6, 7, and 12), five were born in other areas of Southern Greece, and two others were born in Turkey and migrated to Greece during their adolescence. All pursued different professional activities but most of them were still living as farmers sometimes under poor sociohygienic conditions. To the best of the patients' knowledge, none of their closer relatives had overt signs of KS. All patients studied were heterosexuals with no history of sexually transmitted diseases. None of them had ever experienced any severe infectious disease nor had anyone received blood transfusions, therapy with blood products, or any immunosuppressive treatment.

**Clinical and Routine Laboratory Evaluation** In addition to a detailed physical evaluation, patients were subjected to various radiologic, sonographic, and endoscopic procedures. Routine investigations included blood sedimentation rate, complete and differential blood cell counts, serum electrolytes, renal and liver function tests, and syphilis serology (VDRL, TPHA tests).

**Virus Serology** Cytomegalovirus (CMV), herpes simplex virus (HSV), and varicella zoster virus (VZV)-specific immunoglobulin (Ig)M and IgG antibodies were determined by solid-phase enzyme-linked immunosorbent assays (ELISA) according to established procedures [14,15]. Antibodies against the Epstein-Barr virus (EBV) capsid antigen (VCA), EBV early antigens (EA) and EBV nuclear antigens (EBNA) were determined by indirect immunofluorescence (IF) assays [16,17]. The test for human herpes virus type 6 (HHV-6)-specific IgG antibodies was carried out using the HHV-6 kit from Pan-Data-Systems, Inc. (Rockville, MD).

Seroreactivity testing against human retroviruses was performed in two separate laboratories and included the screening for serum

antibodies against (a) HIV-1 by commercial ELISA (Organon Teknika, Turnhout, Belgium) and Western blot (Dupont-Comp. Med. Prod., Wilmington, DE) assays; and against (b) HIV-2 and (c) human T-lymphotropic virus 1 (HTLV-1) by radioimmunoprecipitation assays using appropriate virus-infected cell lines [18,19].

**Immunomorphologic Studies on Biopsies from KS Skin Lesions** One to three biopsies of KS lesions in different stages of development were taken from each patient from different body regions under local anesthesia with 2% mepivacain; each biopsy was divided into three parts: one part was immediately fixed in 1.96% paraformaldehyde, pH 7.4, and further processed for histopathological examination; a second portion of the specimen was fixed in a mixture of 2% glutaraldehyde and 2% paraformaldehyde in 0.1 M cacodylate buffer, pH 7.4, and then further processed for transmission electronmicroscopic analysis.

For immunohistologic studies, the third part of the biopsy specimen was snap-frozen in Tissue-Tek OCT compound (Miles Scientific, Naperville, IL) using isopentane pre-cooled in liquid nitrogen and then stored at  $-70^{\circ}\text{C}$  until further use. Four-micrometer cryostat serial sections were mounted onto gelatine-coated slides, air-dried, and fixed in acetone for 10 min at  $4^{\circ}\text{C}$ . For blocking of Fc-Ig receptors, sections were preincubated for 20–30 min at room temperature (RT) in phosphate-buffered saline (PBS, pH 7.4) containing 20% heat-inactivated sheep serum (GIBCO, Grand Island, NY). For morphologic orientation, the first cryostat section of each tissue block was stained with hematoxylin and eosin.

First-step antibodies used are listed in Table I [20–34].

**Immunoenzymatic Staining** For single-color staining, the three-step avidin-biotin immunoperoxidase technique was used as described elsewhere [35]. Briefly, sections were incubated with the first-step antibodies (working dilution, see Table I) for 12–16 h at  $4^{\circ}\text{C}$ . Thereafter, the slides were thoroughly washed in PBS, and then reacted with either an appropriately diluted biotin-conjugated sheep anti-mouse Ig (Amersham, Amersham, UK) or a biotin-conjugated donkey anti-rabbit Ig (Amersham) for 1 h at RT. After three PBS washes, slides were further incubated with the streptavidin-biotin peroxidase complex (Strept ABCComplex-HRP, Dakopatts; Copenhagen, Denmark) as described in the data sheet. Bound immunoreactants were visualized by using the diaminobenzidine reaction.

For double-labeling purposes, the incubation chain was extended by consecutively exposing the sections to appropriate murine MoAb, biotin-labeled sheep anti-mouse Ig and, finally, avidin-biotin alkaline phosphatase complex (ABComplex-AP, Dakopatts). This enzyme was then visualized with naphthol AS-MX phosphate/Fast Red [35].

**Controls** Cryostat sections of skin (healthy volunteers) and lymph node specimens (autopsy material) were used to define the optimal working dilutions of the respective antibodies.

Negative controls included (a) omission of either the first or one

**Table III.** Immunophenotype of KS Cell<sup>a</sup>

MoAb	Normal Human Skin		KS Lesions	
	Endothelium of Mature Blood Vessels/Lymphatic		Endothelium of Clefts/Slits	Spindle Cells
PAL-E	++	—	—	—
EN-4	++	++	+	+
Anti-von Willebrand factor	+	—	—	—
OKM-5 (anti-CD36)	+	—	—/(+)	—/(+)
Anti-laminin	++	+	+	+
Anti-collagen type-4	++	+	+	+
Anti-desmin	—	—	—	—
Anti-vimentin	++	++	++	++

<sup>a</sup>—, negative; (+), occasionally weakly positive; +, weakly positive; ++, strongly positive.

of the subsequent antibody reagents; (b) substitution of the primary antibody either with normal mouse or rabbit serum; (c) substitution of the first antibody with an irrelevant antibody of the same isotype. Consistently, negative results were obtained.

## RESULTS

**Clinical Features** Each of the 12 patients displayed multiple KS lesions in all stages of development (Fig 1). The lesions were most pronounced on the distal portions of the lower and upper extremities but also occurred, at varying frequency, on the proximal parts of the extremities, head, neck, and face. Characteristically, the lesions were predominantly located on acral areas and were usually distributed in a symmetrical fashion (Fig 1, Table II). In addition to the cutaneous lesions, mucosal involvement was observed in 10 patients. Four patients exhibited orogenital KS patches; endoscopic evaluation of the gastrointestinal tract revealed esophageal, gastric, and/or duodenal KS lesions in nine of 10 patients examined (Table II). In one third of the patients, indolent palpable lymph nodes (diameter approximately 2 cm) were noted in the inguinal (patients 3, 4, and 6) and cervical (patients 1 and 6) regions. Unfortunately, lymph node biopsies for histologic examination were not feasible. With the exception of patient 4, who had experienced a weight loss of 10 kg during the last 4 months prior to examination, the patients were in good general health and were not incapacitated by the KS lesions. None of the patients offered a history of overt melena, and radiologic and/or sonographic studies of thorax and abdomen did not reveal any pathologic process.

**Laboratory Data** The routine laboratory data (blood sedimentation rate, total and differential blood cell count, electrolytes, liver and renal function tests) were essentially within normal limits. Syphilis serology tests were negative in all patients.

Antibodies to HTLV-1, HIV-1, and HIV-2 were not detectable in any of the patients' sera by either ELISA, Western blot, or radioimmunoprecipitation assays. Each patient, however, exhibited detectable IgG antibody titers to HSV, VZV, CMV, HHV-6, and EBV. In the latter instance, the antibody spectrum included IgA antibodies against EBV-VCA and antibodies against EBNA and EBV-EA.

**Histopathology** The histopathologic examination of biopsy specimens revealed the typical picture of the various stages of KS. The early patch-like lesions predominantly consisted of increased numbers of irregularly sized, jagged-shaped, thin-walled clefts and slits, dissecting the pre-existing dermal collagen bundles (Fig 2a). Erythrocytes were frequently detected within these clefts but were occasionally also extravasated in the spongy network of collagen fibers. Elongated cells lining these clefts did not show any signs of atypia and were considered to be endothelial cells. In addition to the changes already seen in the patch stage, the advanced plaque stage showed some vascular neoproliferation consisting of normal-appearing venules and arterioles; a distinct sign of the plaque lesions were solid cords composed of spindle cells arranged in an indian file pattern. This biphasic tumor morphology, exhibiting both angiomatous and solid tumor pattern, changed to a clearcut sarcomatous

pattern in the progressed nodular stage. These lesions consisted almost exclusively of spindle cells that were arranged in bundles and interlacing fascicles (Fig 2b). Some of the tumors were relatively sharply demarcated, exhibiting a pseudocapsule at the periphery; others were characterized by an invasive growth throughout the entire reticular dermis. In high magnification only, we observed numerous erythrocytes between the individual spindle cells as well as siderophages that were predominantly found at the periphery of the tumors. Signs of pronounced cellular atypia or an increased mitotic rate were rarely seen in any stages of development. A hallmark of all lesions was the admixture of a moderate to dense inflammatory infiltrate, consisting of lymphocytes, histiocytes, and plasma cells (see below).

## Immunohistochemistry


**Phenotypic Analysis of the Vascular Component of KS Lesions** In biopsies of normal human skin from healthy volunteers, we observed that MoAb EN-4 and anti-factor VIII-related antigen (anti-von Willebrand Factor (vWF)) displayed selective reactivity with endothelial cells. Clearcut endothelial cell reactivity was also seen with MoAb PAL-E and OKM-5, but these reagents also reacted with other structures/cells (Table III). In agreement with previous observations, we found (a) that MoAb EN-4 reacted with both lymphatic and blood vessel endothelium; (b) that PAL-E, anti-vWF, and OKM-5 reacted with blood vessel endothelial cells only; and (c) that all endothelial cells expressed vimentin, but failed to react with anti-desmin antibodies (Table III) [21–29].

In all specimens of patch/plaque stage KS lesions, cells lining the typical clefts and slits were uniformly EN-4-positive, but virtually failed to display PAL-E, OKM-5, and anti-vWF reactivity (Table III). At their abluminal side, these cells exhibited distinct anti-laminin and anti-type IV collagen reactivity indicative of the presence of a basal lamina. Thus, the phenotypic profile of these cells is indistinguishable from that of lymphatic endothelial cells in normal human skin (Table III).


In late plaque and tumor-stage KS lesions, spindle cells forming solid tumor nests and bundles were again EN-4+, PAL-E-, and anti-vWF- and only occasionally displayed weak OKM-5 reactivity (Table III, Fig 3). Remarkably, these regions showed a weak but distinct anti-laminin- and anti-collagen type IV staining. Although this immunolabel could not be ascribed to defined structures, it clearly coincided with tumorous areas and did not exceed their outer contours. Similar to the situation in normal human skin, anti-collagen type IV and anti-laminin staining outside the proliferating spindle cells were confined to basement membranes of both epithelial-dermal junctions and blood vessels. In contrast to the cells lining the pathognomonic vascular channels mentioned above, endothelial cells lining these blood vessels showed strong PAL-E, EN-4, anti-vWF, and moderate OKM-5 immunolabeling (Table III, Fig 3). Finally, it should be emphasized that the spindle cells as well as the endothelial cells were uniformly of the vimentin rather than of the desmin type.

**Phenotypic Analysis of Inflammatory Cells Within KS Lesions** Immunolabeling of serial sections revealed that EN-4+ cells were lack-





**Figure 3.** Immunophenotypic analysis of a solid KS lesion on serial-cryostat sections. *a*, mature blood vessels (arrowheads) as well as spindle cells (arrows) are labeled with MoAb EN-4. *b*, MoAb PAL-E stains endothelial cells of mature blood vessels (arrowheads) but fails to react with spindle cells. *c*, labeling with anti-vWF MoAb is restricted to mature blood vessels (arrowheads); the spindle cells are negative.



**Figure 4.** Immunophenotypic analysis of mononuclear cells infiltrating KS lesions. CD4<sup>+</sup> cells (brownish reaction product) greatly outnumber CD8<sup>+</sup> lymphocytes (red membrane staining.).


ing the common leukocyte antigen CD45. However, considerable numbers of CD45<sup>+</sup> cells were seen around patch-stage vascular slits as well as between and in the immediate vicinity of spindle cell cords, nests, and bundles. Most of these cells represented CD3<sup>+</sup> lymphocytes which, by appropriate serial sectioning, were predominantly of the CD4 subtype (Fig 4). To a lesser extent, we also identified CD14-bearing macrophages and anti-factor XIIIa-reactive stellate and elongated cells [34]. CD20<sup>+</sup> B cells were barely detectable.

### Electronmicroscopy

**Cell Morphology** The irregularly shaped clefts and slit-like spaces of patch and plaque-stage KS lesions were outlined by a monolayer of cells that appeared rather thin with elongated processes, sometimes protruding into the surrounding collagen tissue. The cells formed intercellular connections (tight junctions, desmosome-like structures, overlapping and interdigitating junctions), exhibited large numbers of pinocytotic vesicles and multivesicular bodies but were consistently devoid of Weibel-Palade bodies. At the abluminal side, the cells were inconsistently outlined by a basal lamina but pericytes were not observed (Fig 5a). Thus, these cells greatly resemble lymphatic endothelial cells. Most noteworthy was the additional finding of tubulo-reticular structures (TRS, Fig 5a), erythrophagocytosis, and membrane ghosts within their cytoplasm.

Spindle cells comprising fascicles displayed similar morphologic features including the presence of intercellular junctions, pinocytotic vesicles, multivesicular bodies, and TRS, but, in addition, also occasionally contained cylindrical confronting cisternae (CCC) (Fig 5b). A careful screening (more than 3000 tumor cells examined) for the presence of Weibel-Palade bodies yielded negative results with the exception of two spindle cells each of which contained one single Weibel-Palade body. Most of the spindle cells were in tight contact with each other and, thus, no longer associated with vascular lumina and not surrounded by a basal lamina. Occasionally, however, widened channel-like spaces were observed between neighboring spindle cells and in these instances basal lamina formation was observed at the "abluminal" side (Fig 5c).





**Figure 5.** Ultrastructural characteristics of KS cells. *a*, TRS (arrow) within a cell forming the endothelium of a slit-like lumen. Note the basal lamina at the abluminal side (arrowheads) and its fragmentation (asterisks). No Weibel-Palade bodies can be observed. L, lumen; magnification  $\times 13,000$ ; bar,  $1\ \mu\text{m}$ . *b*, CCC (arrowheads) observed in a spindle cell of a KS lesion; magnification  $\times 30,000$ ; bar,  $500\ \text{nm}$ . *c*, spindle cells form abortive vessel-like channels (arrowheads) with multiple pinocytotic vesicles (small arrows); at the "abluminal side" remnants of a basal lamina can be discerned (bold arrows); magnification  $\times 24,000$ ; bars,  $500\ \text{nm}$ .

As opposed to these spindle cells, morphologically normal-appearing blood vessels were lined by cells displaying the typical features of blood vessel endothelial cells, i.e., prominent nuclei and the almost regular presence of Weibel-Palade bodies. These blood vessels were surrounded by a continuous, often multilayered basement membrane and adjoining pericytes.

**Retrovirus-Like Particles** Extensive electronmicroscopic tissue screening revealed the presence of virus-like particles in biopsy specimens of KS lesions from five patients (patients 3, 5, 8, 11, and 12). These round/globular structures measured  $90\text{--}130\ \text{nm}$  in overall diameter. Some of these particles consisted of a ring-shaped, electron-dense nucleoid ( $60\text{--}70\ \text{nm}$  in diameter) and a more electron-lucent center. The ring-shaped nucleoid was surrounded by a slightly electron-transparent intermediate layer that was enveloped by a peripheral unit membrane (Fig 6a). Multiple surface knobs with a length of  $7\text{--}10\ \text{nm}$  protruded from this membrane (Fig 6a). There existed other particles of similar size that contained a condensed nucleoid of  $40\text{--}50\ \text{nm}$  in diameter (Fig 6b–d). This electron-dense core was located either centrally or slightly eccentrically, was surrounded by an electron-lucent layer of  $25\text{--}30\ \text{nm}$  thickness, and was enveloped by a peripheral unit membrane (Fig 6b–d). On the surface of this membrane, we occasionally detected truncated, but never full-sized, knobs (Fig 6b–d). In essence, the former particles correspond in size and shape to immature retroviral particles and the latter are morphologically indistinguishable from mature oncovirinae [36–38]. These particles were only detected in close vicinity to or, occasionally, even on the surface of tumor cells (Fig 6b–d). In one instance, we detected a mature retroviral particle within a coated pit of a spindle cell (Fig 6e) indicative of its subsequent uptake. In keeping with this assumption was our additional finding of a similar retroviral particle within a cytoplasmic vacuole of an endothelial cell (Fig 6f). Finally, and most importantly, we observed a particle exhibiting the above-described morphologic fea-

tures budding from the membrane of a cytoplasmic vacuole of a tumor cell (Fig 6g).


## DISCUSSION

In this report, we present data on a newly recognized endemic occurrence of KS in a defined Mediterranean population. Epidemiologically, this Greek KS variant resembles the endemic KS form occurring in central Africa that was first documented during the 1950s and 1960s [3,4]. The incidence of eight new cases per year per million Peloponnesian inhabitants refers to the entire peninsula. If one focuses on the geographically more restricted regions of the Southern Peloponnesus (Laconia, Messinia), the incidence is 3 to 4 times higher, and thus approximates the rate of occurrence of the African variant [3,4]. It should also be emphasized that the proportion of affected women is exceedingly high when compared to classical and epidemic KS forms [1,2].

The patients in this study presented a combination of the characteristic clinical features of the different KS variants. Although KS lesions in several stages of development on the feet and hands are representative of classic KS and the "nodular"-type of the African KS, widespread, coin-sized, and streaky lesions on the arms and legs are more commonly seen in patients suffering from the acquired immunodeficiency syndrome (AIDS) [1,2,12]. Moreover, the extensive involvement of the facial skin and head, of the oral and genital mucosa and, more importantly, of the gastrointestinal tract is uncommon in the classic, more common in the African, but a typical feature of the AIDS-associated KS [1,2,12].

Confirming earlier histopathologic reports [4,6,12,39], the initial patch-like lesions consisted of increased numbers of bizarre, endothelium-lined clefts and slits which were accompanied by solid tumor cords consisting of spindle cells in the plaque stage and acquiring a sarcomatous appearance in the nodular stage. In accordance with previous observations, we found that cells lining cleft-like





**Figure 6.** Identification of retroviral particles in KS lesions. *a*, immature extracellular retrovirus. Note the 7–10-nm long spikes/knobs at the enveloping membrane (thin arrows) and the ring-shaped nucleoid (bold arrows); magnification  $\times 140,000$ . *b–d*: several oncovirinae-like structures are found extracellularly, in close vicinity to the cytoplasmic membrane of KS cells. Note the roundish, centrally or slightly eccentrically located nucleoid; magnification  $\times 80,000$  (*b*),  $\times 100,000$  (*c*),  $\times 125,000$  (*d*). *e*, demonstration of a virus-like particle (large arrows) in close vicinity to a coated pit (arrowheads). Note the tail-like structure protruding from the envelope of the virus (small arrow); magnification  $\times 100,000$ . *f*, virus-like particle (arrows) within a cytoplasmic vacuole of a KS cell. (L, lumen, magnification  $\times 52,000$ ; *g*, budding of an immature virion from the membrane of a cytoplasmic vacuole of a KS cell; magnification  $\times 100,000$ . Bars, 100 nm.



lumina as well as the spindle cells comprising tumor bundles have essentially the same phenotype (EN-4<sup>+</sup>, PAL-E<sup>-</sup>, anti-vWF<sup>-</sup>, OKM5<sup>-</sup>/+ [40–42]) and similar ultrastructural features (lack of Weibel-Palade bodies, formation of a discontinuous basal lamina, absence of pericytes [43]) as lymphatic endothelial cells [44]. These findings lead us to the conclusion that the neoplastic vascular cells and the spindle cells represent distinct differentiation stages of the same cell type that we will refer to as KS cell. It remains to be seen whether the KS cell is a true neoplasm of mature lymphatic endothelial cells (as one may predict on the basis of its phenotype) or, less likely, originates from any other member of the endothelial cell family.

Ultrastructural studies revealed another major feature of Greek KS, i.e., the presence of TRS and CCC within the cytoplasm of the proliferating KS cells. So far, TRS and CCC have been described as prominent ultrastructural features in the context of AIDS-KS [45–47]. This phenomenon is apparently independent of KS per se in that these structures occur in a wide variety of cells of HIV-1-infected persons, and their presence within peripheral mononuclear cells was even considered as a diagnostic and prognostic marker of AIDS [48–50]. However, the occurrence of TRS and CCC is not pathognomonic for HIV-1 infection. It is actually a not uncommon sequela of certain viral infections and/or virus-induced neoplasms in different species (reviewed in [51]). The more recent demonstration that these structures can be induced in vitro by interferon- $\alpha/\beta$  [50–52] supports this assumption and therefore suggests that an underlying viral infection is also responsible for the occurrence of TRS/CCC in our patients.

Indeed, extensive electronmicroscopic sampling of KS lesions revealed the presence of rounded 110–130 nm particles exhibiting an electron-dense central core surrounded by an enveloping unit membrane; in addition, we found a particle of similar size with a ring-shaped electron-dense nucleoid and an enveloping membrane with surface spikes. Thus, these particles fulfilled the established morphologic criteria for both mature and immature retroviruses [36–38]. The assumption that the particles disclosed in our study are true virions infecting KS cells rather than some ill-defined “look alike” structures is further supported, albeit not definitively proved, by the detection of such particles exclusively in close association with KS cells in five of 12 patients and by the unequivocal demonstration of a budding particle in a KS cell in one patient.

We are not aware of reports documenting the *in situ* occurrence of retroviral particles in conjunction with human (tumor) cells—with the notable exception that similar particles have been found budding from KS cells of AIDS patients [53,54]. Although our search for HIV-1, HIV-2, and HTLV-I did not include immunohistologic staining for viral proteins, virus culture, or viral gene amplification by the polymerase chain reaction, our repeated attempts to serologically detect antibodies against proteins encoded by these viruses consistently yielded negative results. This makes it very unlikely that our patients are infected with any of the known human retroviruses. Thus, the possibility should be entertained that the retroviral particles detected in KS lesions from Greek individuals (this study) and, possibly, from AIDS patients [53,54] represent an as yet unidentified human retrovirus.

The crucial question remains whether these newly discovered retroviral particles are of any pathogenetic significance. It is conceivable that they represent exogenous or endogenous non-pathogenic retroviruses occurring exclusively in KS patients. Alternatively, it might even be that extensive ultrastructural tissue screening would reveal their presence in normal skin from healthy individuals.

Should these retroviral particles be involved in KS tumorigenesis, they might trigger events directly resulting in the transformation of endothelial cells. Indeed, others have shown that KS-DNA can transform NIH 3T3 cells [55,56] and that such transformants can induce the appearance of hemorrhagic angiosarcomatous neoplasms in nude mice [56]. On the other hand, the role of retroviruses in KS development could be to induce the production/secretion of angiogenic factors in a given cell as has been reported for CD4<sup>+</sup> T cells

infected with human retroviruses [57]. Such factors may then act in either a paracrine or autocrine fashion to promote/enhance neoplastic proliferation [57–60]. It remains to be seen whether similar interactions are operative between infiltrating CD4<sup>+</sup> T cells and KS cells in our patients.

Our laboratory is currently attempting to rescue and molecularly characterize this putative new retrovirus. The success of these efforts is certainly a *conditio sine qua non* for further studies to probe a possible biologic role for this retrovirus.

*We are indebted to Dr. Hans Gelderblom (Robert Koch-Institut, West Berlin, FRG) for critically reading the manuscript and for helpful advice concerning the virologic aspects of this study. We also thank Mr. Heinz Gaudek, Ms. Ursula Hornick, and Ms. Inge Wadlig for excellent technical assistance, Mrs. Renate Kopp and Sabine Seizov for carefully typing the manuscript, and Mrs. Laura Stingl for helpful suggestions in its preparation.*

## REFERENCES

1. Ziegler JL, Dorfman RF (eds.): Kaposi's sarcoma. Pathophysiology and clinical management. Marcel Dekker, Inc., New York and Basel, 1988
2. Safai B: Kaposi's sarcoma: an overview of classical and epidemic forms. In: Broder S (ed.). AIDS. Modern concepts and therapeutic challenges. Marcel Dekker, Inc., New York and Basel, 1987, pp 205–218
3. Slavin G, Cameron HMcD, Singh H: Kaposi's sarcoma in mainland Tanzania: a report of 117 cases. *Br J Cancer* 23:349–357, 1969
4. Taylor JF, Templeton AC, Vogel CL, Ziegler JL, Kyalwazi SK: Kaposi's sarcoma in Uganda: a clinico-pathological study. *Int J Cancer* 8:122–135, 1971
5. Penn I: Kaposi's sarcoma in organ transplant recipients: report of 20 cases. *Transplantation* 27:8–11, 1979
6. Akhtar M, Bunuan H, Ali MA, Godwin JT: Kaposi's sarcoma in renal transplant recipients. Ultrastructural and immunoperoxidase study of four cases. *Cancer* 53:258–266, 1984
7. Shmueli D, Shapira Z, Yussim A, Nakache R, Ram Z, Shaharabani E: The incidence of Kaposi sarcoma in renal transplant patients and its relation to immunosuppression. *Transplant Proc* 21:3209–3210, 1989
8. Hymes KB, Cheung T, Greene JB, Prose NS, Marcus A, Ballard H, William DC, Laubenstein LJ: Kaposi's sarcoma in homosexual men—a report of eight cases. *Lancet* II:598–600, 1981
9. Gottlieb MS, Schroff R, Schanker HM, Weisman JD, Fan PT, Wolf RA, Saxon A: Pneumocystis carinii pneumonia and mucosal candidiasis in previously healthy homosexual men. Evidence of a new acquired cellular immunodeficiency. *N Engl J Med* 305:1425–1431, 1981
10. Masur H, Michelis MA, Greene JB, Onorato I, Vande Stouwe RA, Holzman RS, Wormser G, Brettman L, Lange M, Murray HW, Cunningham-Rundles S: An outbreak of community-acquired pneumocystis carinii pneumonia. Initial manifestation of cellular immune dysfunction. *N Engl J Med* 305:1431–1438, 1981
11. Siegal FP, Lopez C, Hammer GS, Brown AE, Kornfeld SJ, Gold J, Hassett J, Hirschman SZ, Cunningham-Rundles C, Adelsberg BR, Parham DM, Siegal M, Cunningham-Rundles S, Armstrong D: Severe acquired immunodeficiency in male homosexuals, manifested by chronic perianal ulcerative herpes simplex lesions. *N Engl J Med* 305:1439–1444, 1981
12. Friedman-Kien AE, Laubenstein LJ, Rubinstein P, Buimovici-Klein E, Marmor M, Stahl R, Spigland I, Kim KS, Zolla-Pazner S: Disseminated Kaposi's sarcoma in homosexual men. *Ann Intern Med* 96:693–700, 1982
13. Kaloterakis A: Kaposi's sarcoma in Greece. (Mediterranean Kaposi's sarcoma). Thesis, Department of General Medicine, University of Athens, 1984
14. Stagno S, Reynolds DW, Smith RJ: Use of isolated nuclei in the indirect fluorescent-antibody test for human cytomegalovirus infection: comparison with microneutralisation, anticomplement and conventional indirect fluorescent-antibody assays. *J Clin Microbiol* 7:486–489, 1978



15. Cremer NE, Cossen CK, Hanson CV, Shell GR: Evaluation and reporting of enzyme immunoassay determinations of antibody to herpes simplex virus in sera and CSF. *J Clin Microbiol* 15:815-823, 1982
16. Henle W, Henle G, Horwitz CA: Epstein-Barr virus-specific diagnostic tests in infectious mononucleosis. *Hum Pathol* 5:551-565, 1974
17. Nikoskelainen J, Ablashi DV, Isenberg RA, Neel EV, Miller RG, Stevens DA: Cellular immunity in infectious mononucleosis. II. Specific reactivity to Epstein-Barr virus antigens and correlation with clinical and hematological parameters. *J Immunol* 121:1239-1244, 1978
18. Zagury JF, Franchini G, Reitz M, Collalti E, Starcich B, Hall L, Fargnoli K, Jagodzinski L, Guo HG, Laure F, Arya SK, Josephs S, Zagury D, Wong-Staal F, Gallo RC: Genetic variability between isolates of human immunodeficiency virus (HIV) type 2 is comparable to the variability among HIV type 1. *Proc Natl Acad Sci USA* 85:5941-5945, 1988
19. Popovic M, Sarin PS, Robert-Gurroff M, Kalyanaraman VS, Mann D, Minowada J, Gallo RC: Isolation and transmission of human retrovirus (human T cell leukemia virus). *Science* 219:856-859, 1983
20. Schlingemann RO, Dingjan GM, Emeis JJ, Blok J, Warnaar SO, Ruiter DJ: Monoclonal antibody PAL-E specific for endothelium. *Lab Invest* 52:71-76, 1985
21. Holden CA, Spaul J, Williams R, Spry CJF, Russel Jones R, Wilson Jones E: The detection of endothelial cell antigens in cutaneous tissue using methacarn and periodate lysine paraformaldehyde fixation. *J Immunol Methods* 91:45-52, 1986
22. Russel Jones R, Spaul J, Spry C, Wilson Jones E: Histogenesis of Kaposi's sarcoma in patients with and without acquired immune deficiency syndrome (AIDS). *J Clin Pathol* 39:742-749, 1986
23. Beckstead JH, Wood GS, Fletcher V: Evidence for the origin of Kaposi's sarcoma from lymphatic endothelium. *Am J Pathol* 119:294-300, 1985
24. Rutgers JL, Wiczorek R, Bonetti F, Kaplan KL, Posnett DN, Friedman-Kien AE, Knowles II DM: The expression of endothelial cell surface antigens by AIDS-associated Kaposi's sarcoma. Evidence for a vascular endothelial cell origin. *Am J Pathol* 122:493-499, 1986
25. Knowles II DM, Tolidjian B, Marboe C, D'Agati V, Grimes M, Chess L: Monoclonal anti-human monocyte antibodies OKM 1 and OKM 5 possess distinctive tissue distributions including differential reactivity with vascular endothelium. *J Immunol* 132:2170-2173, 1984
26. Barsky SH, Baker A, Siegal GP, Togo S, Liotta LA: Use of anti-basement membrane antibodies to distinguish blood vessel capillaries from lymphatic capillaries. *Am J Surg Pathol* 7:667-677, 1983
27. Kramer RH, Fuh G-M, Hwang CBC, Conant MA, Greenspan JS: Basement membrane and connective tissue proteins in early lesions of Kaposi's sarcoma associated with AIDS. *J Invest Dermatol* 84:516-520, 1985
28. Leu HJ, Odermatt B: Multicentric angiosarcoma (Kaposi's sarcoma). Light and electron microscopic and immunohistological findings of idiopathic cases in Europe and Africa and of cases associated with AIDS. *Virchows Arch (Pathol Anat)* 408:29-41, 1985
29. Schulze H-J, Rütten A, Mahrle G, Steigleder GK: Initial lesions of HIV-related Kaposi's sarcoma—a histological, immunohistochemical, and ultrastructural study. *Arch Derm Res* 279:499-503, 1987
30. Beverley PCL, Linch D, Delia D: Isolation of human haematopoietic progenitor cells using monoclonal antibodies. *Nature* 287:332-333, 1980
31. Ledbetter JA, Evans RL, Lipinski M, Cunningham-Rundles C, Good RA, Herzenberg LA: Evolutionary conservation of surface molecules that distinguish T lymphocyte helper/inducer and cytotoxic/suppressor subpopulations in mouse and man. *J Exp Med* 153:310-323, 1981
32. Dimitriu-Bona A, Burmester GR, Waters SJ, Winchester RJ: Human mononuclear phagocyte differentiation antigens. I. Patterns of antigenic expression on the surface of human monocytes and macrophages defined by monoclonal antibodies. *J Immunol* 130:145-152, 1983
33. Stashenko P, Nadler LM, Hardy R, Schlossman SF: Characterization of a human B lymphocyte-specific antigen. *J Immunol* 125:1678-1685, 1980
34. Nickoloff BJ, Griffiths CEM: Factor XIIIa-expressing dermal dendrocytes in AIDS-associated cutaneous Kaposi's sarcomas. *Science* 243:1736-1737, 1989
35. Gillitzer R, Pilarski LM: In situ localization of CD45 isoforms in the human thymus indicates a medullary location for the thymic generative lineage. *J Immunol* 144:66-74, 1990
36. Bouillant AMP, Becker SAWE: Ultrastructural comparison of oncovirinae (Type C), spumavirinae and lentivirinae: three subfamilies of retroviridae found in farm animals. *J Natl Canc Inst* 72:1075-1084, 1984
37. Munn RJ, Marx PA, Yamamoto JK, Gardner MB: Ultrastructural comparison of the retroviruses associated with human and simian acquired immunodeficiency syndromes. *Lab Invest* 53:194-199, 1985
38. Gelderblom HR, Özel M, Pauli G: Morphogenesis and morphology of HIV structure-function relations. *Arch Virol* 106:1-13, 1989
39. Dorfman RF: Kaposi's sarcoma revisited. *Hum Pathol* 15:1013-1017, 1984
40. Scully PA, Steinman HK, Kennedy C, Trueblood K, Frisman DM, Voland JR: AIDS-related Kaposi's sarcoma displays differential expression of endothelial surface antigens. *Am J Pathol* 130:244-251, 1988
41. Green TL, Meyer JR, Daniels TE, Greenspan JS, de Souza Y, Kramer RH: Kaposi's sarcoma in AIDS: basement membrane and endothelial cell markers in late stage lesions. *J Oral Pathol* 17:266-272, 1988
42. Nadimi H, Saatee S, Armin A, Toto PD: Expression of endothelial cell markers PAL-E and EN-4 and Ia antigens in Kaposi's sarcoma. *J Oral Pathol* 17:416-420, 1988
43. Mc Nutt NS, Fletcher V, Conant MA: Early lesions of Kaposi's sarcoma in homosexual men. An ultrastructural comparison with other vascular proliferations in skin. *Am J Pathol* 111:62-77, 1983
44. Daroczy J (ed.): The dermal lymphatic capillaries. Springer-Verlag, Berlin-Heidelberg, 1988
45. Gyorkey F, Sinkovics JG, Gyorkey P: Tubulo-reticular structures in Kaposi's sarcoma. *Lancet* II:984-985, 1982
46. Sidhu GS, Stahl RE, El-Sadr W, Zolla-Pazner S: Ultrastructural markers of AIDS. *Lancet* I:990-991, 1983
47. Sidhu GS, Stahl RE, El-Sadr W, Cassai ND, Forrester EM, Zolla-Pazner S: The acquired immunodeficiency syndrome: an ultrastructural study. *Hum Pathol* 16:377-386, 1985
48. Orenstein JM, Simon GL, Kessler CM, Schulof RS: Ultrastructural markers in circulating lymphocytes of subjects at risk for AIDS. *Am J Clin Pathol* 84:603-609, 1985
49. Kostianovsky M, Grimley PM: Ultrastructural findings in the acquired immunodeficiency syndrome (AIDS). *Ultrastruct Pathol* 8:123-130, 1985
50. Yoffe B, Petrie BL, Noonan CA, Hollinger FB: In vivo and in vitro ultrastructural alterations induced by human immunodeficiency virus in human lymphoid cells. *Lab Invest* 61:303-309, 1989
51. Luu J, Bockus D, Remington F, Bean MA, Hammar SP: Tubuloreticular structures and cylindrical confronting cisternae: a review. *Hum Pathol* 20:617-627, 1989
52. Orenstein JM, Preble OT, Kind P, Schulof R: The relationship of serum alpha-interferon and ultrastructural markers in HIV-seropositive individuals. *Ultrastruct Pathol* 11:673-679, 1987
53. Gyorkey F, Sinkovics JG, Melnick JL, Gyorkey P: Retroviruses in Kaposi-sarcoma cells in AIDS. *N Engl J Med* 311:1183-1184, 1984
54. Schenk P, Konrad K: Ultrastructure of Kaposi's sarcoma in acquired immune deficiency syndrome (AIDS). *Arch Otorhinolaryngol* 242:305-313, 1985
55. Delli-Bovi P, Basilico C: Isolation of a rearranged human transforming gene following transfection of Kaposi sarcoma DNA. *Proc Natl Acad Sci USA* 84:5660-5664, 1987
56. Lo SC, Liotta LA: Vascular tumors produced by NIH/3T3 cells transfected with human AIDS Kaposi's sarcoma DNA. *Am J Pathol* 118:7-13, 1985
57. Nakamura S, Salahuddin SZ, Biberfeld P, Ensoli B, Markham PD, Wong-Staal F, Gallo RC: Kaposi's sarcoma cells: long-term culture with growth factor from retrovirus-infected CD4<sup>+</sup> cells. *Science* 242:426-430, 1988



58. Salahuddin SZ, Nakamura S, Biberfeld P, Kaplan MH, Markham PD, Larsson L, Gallo RC: Angiogenic properties of Kaposi's sarcoma-derived cells after long-term culture in vitro. *Science* 242:430-433, 1988
59. Ensoli B, Nakamura S, Salahuddin SZ, Biberfeld P, Larsson L, Beaver B, Wong-Staal F, Gallo RC: AIDS-Kaposi's sarcoma-derived cells express cytokines with autocrine and paracrine growth effects. *Science* 243:223-226, 1989
60. Ensoli B, Barillari G, Salahuddin SZ, Gallo RC, Wong-Staal F: Tat protein of HIV-1 stimulates growth of cells derived from Kaposi's sarcoma lesions of AIDS patients. *Nature* 345:84-86, 1990

## ANNOUNCEMENT

We are pleased to announce that the 1993 Tricontinental Meeting of The European Society for Dermatologic Research, The Japanese Society for Investigative Dermatology, and The Society for Investigative Dermatology will be held in Japan in the Fall of 1993. We expect that there will be a limited number of grants available to qualified junior investigators, residents, and fellows to help offset travel costs. In addition, the societies are exploring the possibility of obtaining funding to support visiting professorships in Japan at the time of the meeting. Further information will be published in the *Journal of Investigative Dermatology* and can be obtained by contacting The Society for Investigative Dermatology at (216) 844-3682.